



QUESTIONS to EVALUATE YOUR CHILD'S PENICILLIN or PENICILLIN-LIKE ALLERGY

You have stated that your child has a penicillin or penicillin-like allergy. It is now recommended by the American Academy of Allergy, Asthma and Immunology that all children with a penicillin or penicillin-like allergy have an allergy evaluation proactively. More than 95% of patients labeled as penicillin or penicillin-like allergic are not actually allergic and 80% of people lose their allergy after 10 years.

This questionnaire will help us determine if your child can safely receive a penicillin or penicillin-like antibiotic.

1. There are 4 sections to determine your child's penicillin allergy type.
2. Please answer each question by choosing YES or NO
3. Name of allergic antibiotic(s) _____ or fill in "Unknown"

Section 1

THIS WILL DETERMINE IF YOUR CHILD HAS A RARE SEVERE ALLERGY.

Step 1

Does your child have a history of ANY of the following reactions after taking an antibiotic? **THE REACTION MAY HAVE OCCURRED ALMOST IMMEDIATELY OR UP TO SEVERAL WEEKS** after taking the antibiotic. Please check "Yes" or "No" (or circle Yes or No if printed out). (Titles below are for health care provider use)

1. SJS/TEN
 - Rash devoping into blisters all over your child's body YES NO
 - Mouth sores YES NO
 - Redness of the eyes YES NO
2. DRESS
 - Rash all over your child's body followed by peeling of the skin YES NO
 - Rash with mouth sores, swollen glands, liver or kidney problems YES NO

- | | | | |
|----|--|------------|-----------|
| 3. | <u>AGEP</u> | | |
| | <ul style="list-style-type: none"> • Rapidly spreading, pin-sized pus-filled, blister-like sores all over your child's body | YES | NO |
| 4. | <u>SSLR</u> | | |
| | <ul style="list-style-type: none"> • Rash and joint pains | YES | NO |
| 5. | <u>VASCULITIS</u> | | |
| | <ul style="list-style-type: none"> • Purple rash with kidney or lung damage | YES | NO |
| 6. | <u>ORGAN SPECIFIC</u> | | |
| | <ul style="list-style-type: none"> • Damage to liver, kidneys or blood count | YES | NO |

If “YES” to any question(s), your child is **SEVERELY ALLERGIC** to penicillin and any of the penicillin-like antibiotics.

Your child should not take any penicillin or penicillin-like antibiotics until you are evaluated by your child's healthcare provider.

****YOU DO NOT HAVE TO ANSWER ANY OTHER QUESTIONS.****

If “NO” go to **Step 2**.

Section 2

THIS WILL DETERMINE IF YOU HAVE A SEVERE IMMEDIATE ALLERGY.

Step 2

Does your child have a history of any of the following reactions after taking an antibiotic?

- | | | | |
|---|---|------------|-----------|
| • | Sudden onset of shortness of breath, wheezing, cough or high-pitched whistling sound while taking a deep breath | YES | NO |
| • | Passed out | YES | NO |
| • | Total body rash with welts that itch | YES | NO |
| • | Swollen lips, tongue or throat tightness | YES | NO |
| • | Abdominal pain or vomiting | YES | NO |

If “YES” to any question(s), your child is **SEVERELY ALLERGIC** to penicillin and any of the penicillin-like antibiotics.

Your child should not take any penicillin or penicillin-like antibiotics until you are evaluated by your child’s healthcare provider.

****YOU DO NOT HAVE TO ANSWER ANY OTHER QUESTIONS.****

If “NO” go to **Step 3**.

Section 3

THIS WILL DETERMINE IF YOUR CHILD IS NOT ALLERGIC.

Step 3

Has your child had any of the following **NON ALLERGIC** reactions to an antibiotic?

- | | | |
|---|------------|-----------|
| • Headache | YES | NO |
| • Diarrhea, nausea, vomiting or abdominal pain | YES | NO |
| • Flat rash without welts | YES | NO |
| • Itching without a rash | YES | NO |
| • Does your child have a family history of penicillin allergy | YES | NO |

If “YES”, your child is **NOT ALLERGIC** to penicillin and any of the penicillin-like antibiotics.

****YOU DO NOT HAVE TO ANSWER ANY OTHER QUESTIONS.****

If “NO” go to **Step 4**.

Section 4

THIS WILL DETERMINE IF YOUR CHILD HAS A DELAYED (NON-SEVERE) ALLERGY AND IS AT A LOW RISK FOR TAKING THE ANTIBIOTIC.

Step 4

Since your child's antibiotic reaction, has your child taken any of the following antibiotics without any reaction?

Ampicillin	YES	NO
Amoxicillin	YES	NO
Amoxicillin/Clavulanate (Augmentin)	YES	NO
Cephalexin (Keflex)	YES	NO
Cefadroxil (Duricef)	YES	NO
Cefuroxime (Ceftin)	YES	NO
Cefaclor (Ceclor)	YES	NO
Cefixime (Suprax)	YES	NO
Ceftriaxone (Rocephin)	YES	NO
Penicillin - oral, injection	YES	NO
Cefdinir (Omnicef)	YES	NO

If "YES", your child is NOT ALLERGIC to penicillin and any of the penicillin-like antibiotics.

****YOU DO NOT HAVE TO ANSWER ANY OTHER QUESTIONS.****

If "NO" or "UNKNOWN" go to Step 5.

Step 5

Has the time since the reaction been **greater than 5 years or unknown?** YES NO

If "YES", your child **most likely does not have** a penicillin or penicillin-like allergy or has outgrown it. You need to check with your child's healthcare provider and should not take this type of antibiotic until your child is evaluated.

****YOU DO NOT HAVE TO ANSWER ANY OTHER QUESTIONS.****

If “NO” or “UNKNOWN” go to **Step 6**.

Step 6

Did the reaction occur **greater than 6 hours** after your child received the antibiotic?

YES

NO

If “YES”, your child **most likely does not have** a penicillin or penicillin-like allergy or has outgrown it. You need to check with your child’s healthcare provider and should not take this type of antibiotic until your child is evaluated.

If “NO” to any question(s), your child is **SEVERELY ALLERGIC** to penicillin and any of the penicillin-like antibiotics

Your child should not take any penicillin or penicillin-like antibiotics until your child is evaluated by your child’s healthcare provider.